

PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Health Care Supervision for Children With Williams Syndrome

Committee on Genetics

Pediatrics 2001;107;1192-1204

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://www.pediatrics.org/cgi/content/full/107/5/1192>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2001 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



AMERICAN ACADEMY OF PEDIATRICS

Committee on Genetics

Health Care Supervision for Children With Williams Syndrome

ABSTRACT. This set of guidelines is designed to assist the pediatrician to care for children with Williams syndrome diagnosed by clinical features and with regional chromosomal microdeletion confirmed by fluorescence in situ hybridization.

ABBREVIATIONS. WS, Williams syndrome; FISH, fluorescence in situ hybridization.

INTRODUCTION

Williams syndrome (WS, also Williams-Beuren syndrome), now recognized to be caused by a microdeletion of chromosome 7, is a multisystem disorder first identified as a distinct clinical entity in 1961.¹ It is present at birth and affects boys and girls equally. As routine genetic amniocentesis does not typically detect chromosome microdeletions, children with WS usually come to the attention of pediatricians during infancy or childhood. Initially thought to be a rare genetic disorder, increased awareness of the clinical features and establishment of a reliable diagnostic test have revealed WS to be one of the more commonly recognized genetic disorders in childhood. Williams syndrome is characterized by dysmorphic facies (100%), cardiovascular disease (most commonly supra-aortic stenosis [80%]), mental retardation (75%), a characteristic cognitive profile (90%), and idiopathic hypercalcemia (15%)²⁻⁵ (Table 1).

The diagnosis historically has been made on the basis of clinical criteria (Fig 1), but recently it has been shown that 99% of patients with WS have a hemizygous submicroscopic deletion of 7q11.23 detectable by fluorescence in situ hybridization (FISH).⁶⁻⁸ Chromosome analysis and the Williams Syndrome Chromosomal Region FISH test are recommended for confirmation of the diagnosis. (A child with the clinical features of WS and a negative FISH result should be referred to a clinical geneticist for further evaluation.) The deleted portion of the chromosome includes the *ELN* gene that codes for the structural protein elastin, an important component of the elastic fibers found in the connective tissue of many organs. The *elastin* deletion explains some of the characteristics of WS, such as some of the facial features, hoarse voice, bladder and bowel diverticula, cardiovascular disease, and orthopedic problems. The pathogenesis of other characteristics,

such as hypercalcemia, mental retardation, and unique personality traits, remains unexplained. One possibility is that the loss of 1 or more genes contiguous to the *ELN* gene contributes to the phenotype.

The pediatrician can use knowledge of the clinical manifestations (Table 1) and natural history of WS to anticipate medical problems and to educate the family. Most children with WS are described as having similar facial features.^{4,9} Although these features are often subtle, they tend to become more distinctive with advancing age. Facial features often include periorbital fullness, short nose with bulbous nasal tip, long philtrum, wide mouth, full lips, and mild micrognathia. Infants have full cheeks and a flat facial profile, whereas older children and adults often have a long narrow face and a long neck.^{10,11} Blue- and green-eyed children with WS have a prominent "starburst" pattern to their irides (stellate iris).¹² Mild prenatal growth deficiency and a postnatal growth rate about 75% of normal are consistently observed features of the condition.^{8,13}

The majority of children with WS have cardiovascular anomalies.^{1,2,4} The most common cardiovascular defect is supra-aortic stenosis, an often progressive condition that may require surgical repair.^{10,11} Peripheral pulmonary artery stenosis is often present in infancy and usually improves over time. Coarctation of the aorta, renal artery stenosis, and systemic hypertension are complications that when present may worsen over time.^{4,11,14,15} Because the elastin protein is an important component of elastic fibers in the arterial wall, any artery may become narrowed.

Idiopathic infantile hypercalcemia is an intriguing feature of WS that can contribute to the presence of extreme irritability, vomiting, constipation, and muscle cramps associated with this condition.^{4,9} Symptomatic hypercalcemia usually resolves during childhood, but lifelong abnormalities of calcium and vitamin D metabolism may persist. Hypercalciuria is common and predisposes to nephrocalcinosis. The cause of the abnormality in calcium metabolism is unknown.

An infant with WS often has difficulty feeding and may be brought for medical care because of gastroesophageal reflux, colic, or failure to thrive.^{4,9,16} Other medical problems include Chiari I malformation, strabismus,¹² hyperopia,¹² chronic otitis media, hypodontia, malocclusion, bowel or bladder diverticula, hernias, joint laxity, joint contractures,¹⁷ kyphosis, lordosis, renal or urinary tract malformations,^{14,15} hypothyroidism, and rectal prolapse.

Children with WS have a unique cognitive and

The recommendations in this statement do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

PEDIATRICS (ISSN 0031 4005). Copyright © 2001 by the American Academy of Pediatrics.

TABLE 1. Medical Problems in Williams Syndrome* by Organ System and Age

Organ System	Incidence (%)	Age		
		Infancy	Childhood	Adult
Ocular and visual				
Esotropia	50	x		
Hyperopia	50		x	x
Auditory				
Chronic otitis media	50	x	x	
Hypersensitivity to sound	90	x	x	x
Dental				
Malocclusion	85		x	x
Microdontia	95		x	x
Cardiovascular				
Any abnormality (total)	80	x	x	x
SVAS	75	x	x	x
SVPS	25	x	x	x
PPS	50	x		
Renal artery stenosis	45	x	x	x
Other arterial stenosis	20		x	x
VSD	10	x		
Hypertension	50		x	x
Genitourinary				
Structural anomaly	20	x	x	x
Enuresis	50		x	
Nephrocalcinosis	<5	x	x	x
Recurrent urinary tract infections	30			x
Gastrointestinal				
Feeding difficulties	70	x	x	
Constipation	40	x	x	x
Colon diverticula	30		x	x
Rectal prolapse	15	x	x	
Integument				
Soft lax skin	90	x	x	x
Inguinal hernia	40	x		
Umbilical hernia	50	x		
Prematurely gray hair	90			x
Musculoskeletal				
Joint hypermobility	90	x	x	
Joint contractures	50	x	x	x
Radioulnar synostosis	20	x	x	x
Kyphosis	20			x
Lordosis	40		x	x
Awkward gait	60		x	x
Calcium				
Hypercalcemia	15	x		x
Hypercalciuria	30	x	x	x
Endocrine				
Hypothyroidism	2	x	x	x
Early puberty (but rarely true precocious puberty)	50		x	
Diabetes mellitus	15			x
Obesity	30			x
Neurologic				
Hyperactive deep tendon reflexes	75		x	x
Chiari I malformation	10	x	x	x
Hypotonia (central)	80	x	x	
Hypertonia (peripheral)	50		x	x
Cognitive				
Developmental delay	95	x	x	
Mental retardation	75		x	x
Borderline intellectual functioning	20		x	x
Normal intelligence	5		x	x
Impaired visuospatial constructive cognition	95		x	x
Behavioral				
Attention-deficit hyperactivity disorder	70		x	
Generalized anxiety disorder	80		x	x

* Percentages based on the following: 1) review of rates of complications in several reports of series of patients with Williams syndrome, and 2) database of 315 children and adults with Williams syndrome evaluated by Colleen A. Morris, MD. SVAS indicates supravalvular aortic stenosis; SVPS, supravalvular pulmonic stenosis; PPS, peripheral pulmonary artery stenosis; and VSD, ventricular septal defect.

behavioral profile.^{3,5,18} Cognitive, motor, and language delay are universal, and in 75% of the children, mental retardation is ultimately diagnosed.^{19,20} Older children demonstrate a relative strength in language and auditory memory, with a significant

weakness in visuospatial cognition.^{5,18} Behavioral problems may include hypersensitivity to sound, sleep problems, attention-deficit/hyperactivity disorder,²⁰ and anxiety. Overfriendliness and an empathetic nature are commonly observed.¹⁷

Growth (Past or Present Evidence of) *If 3 of 5 items are checked, score 1 point*

<input type="checkbox"/> Post-term birth > 41 wk gestation	<input type="checkbox"/> Prolonged colic > 4 m irritability
<input type="checkbox"/> Failure to thrive/height and weight < 5th percentile	<input type="checkbox"/> Chronic constipation
<input type="checkbox"/> Vomiting or gastroesophageal reflux	

Behavior and Development

If 3 of 6 items are checked, score 1 point

<input type="checkbox"/> Overly friendly personality	<input type="checkbox"/> Visuospatial problems
<input type="checkbox"/> Hypersensitivity to sound	<input type="checkbox"/> Delayed speech acquisition, followed by excessive talking
<input type="checkbox"/> Anxiety	
<input type="checkbox"/> Developmental delay or mental retardation	

Facial Features

If 8 of 17 items are checked, score 3 points

<input type="checkbox"/> Bitemporal narrowing	<input type="checkbox"/> Broad brow
<input type="checkbox"/> Epicanthal folds or flat nasal bridge	<input type="checkbox"/> Periorbital fullness
<input type="checkbox"/> Strabismus (present or past)	<input type="checkbox"/> Stellate lacy iris pattern
<input type="checkbox"/> Short nose or anteversion of nares	<input type="checkbox"/> Bulbous or full nasal tip
<input type="checkbox"/> Full cheeks	<input type="checkbox"/> Malar hypoplasia (flat cheek bones)
<input type="checkbox"/> Long philtrum	<input type="checkbox"/> Full prominent lips
<input type="checkbox"/> Small, widely spaced teeth	<input type="checkbox"/> Malocclusion
<input type="checkbox"/> Wide mouth	<input type="checkbox"/> Small jaw
<input type="checkbox"/> Prominent ear lobes	

Cardiovascular Problems

(by Echocardiography) (a)

If 1 of 2 items are checked, score 5 points

<input type="checkbox"/> SVAS [†]	<input type="checkbox"/> Peripheral pulmonary artery stenosis
--	---

Cardiovascular Problems (b)

If 1 of 3 items are checked, score 1 point

<input type="checkbox"/> Other congenital heart disease	<input type="checkbox"/> Hypertension
<input type="checkbox"/> Cardiac murmur	

Connective Tissue Abnormality

If 2 of 6 items are checked, score 2 points

<input type="checkbox"/> Hoarse voice	<input type="checkbox"/> Long neck or sloped shoulders
<input type="checkbox"/> Inguinal hernia	<input type="checkbox"/> Joint limitation or laxity
<input type="checkbox"/> Bowel or bladder diverticula	<input type="checkbox"/> Rectal prolapse

Calcium Studies

If 1 of 2 items are checked, score 2 points

<input type="checkbox"/> Hypercalcemia	<input type="checkbox"/> Hypercalciuria
--	---

Total Points:

* If the score is < 3, a diagnosis of Williams syndrome is unlikely. If the score is ≥ 3, FISH studies should be considered. (Mean score for Williams syndrome was 9 [standard deviation = 2.86]. The scoring system is based on a study of 107 persons with Williams syndrome [confirmed by FISH] evaluated by Colleen A. Morris, MD; Frank Greenberg, MD; Paige Kaplan, MD; Martin Levinson, MD; and Barbara Pober, MD; with data analysis by Carolyn B. Mervis, PhD and Byron F. Robinson, MA; presented at the 1994 Williams Syndrome Association Convention; July 31, 1994; San Diego, CA.)

[†] If supralvalvar aortic stenosis (SVAS) is present, referral to a geneticist and FISH studies are recommended.

Fig 1. Williams syndrome diagnostic scoring table: clinical diagnosis.

The medical care of children with WS requires an understanding of the natural history of the disorder, awareness of potential clinical complications, and ongoing assessment and periodic review at appropriate ages (Fig 2). Because the clinical manifesta-

tions during the neonatal period are variable, the diagnosis may not be suspected during early infancy. Accordingly, this statement includes a series of evaluations that should be considered at the time the diagnosis is suspected clinically; the diagnosis

	Neonatal			Infancy (NB - 1 Year)				Early Childhood (1-5 Years)					Late Childhood		Adolescence
		2 mos	4 mos	6 mos	9 mos	12 mos	15 mos	18 mos	24 mos	3 yr	4 yr	5-13 yrs Annual	13-21 yrs Annual		
Diagnosis															
Karyotype/FISH Review [†]	•														
Phenotype Review [†]	•														
Recurrence Risks [†]	•														
Anticipatory Guidance															
Early Intervention	•	•	•	•	•	•	•	•	•						
Family Support	•	•	•	•	•	•	•	•	•	•	•	•	•		
Support Groups [‡]	•	•				•					•	•	•		
Long-term Planning						•					•	•	•		
Sexuality												•	•		
Therapy (pt, ot, speech)										§§		§§	§§		
Medical Evaluation															
Growth feeding	o	o	o	o	o	o	o	o	o	o	o	o	o		
Thyroid Screening	o										o #	o #	o #		
Hearing Screening			s/o			s/o ±			s/o ±		s/o ±	s/o ¶	s/o ¶		
Vision Screening	s/o	s/o	s/o	s/o	s/o ±	s/o			s/o		s/o ±	s/o §	s/o §		
2-Arm Blood Pressure	o			o		o			o	o	o	o	o		
Cardiology Evaluation [†]	**					**			**	†	¶	§	§		
UA/BUN/Cr [†]	o				o	o			o	o	o	o #	o #		
Urine Ca/Cr [†]	o ††					o			o	o	o ¶	o ¶	o ¶		
Serum Calcium [†]	o								o		o	o	o		
Renal Ultrasonography [‡]	o														
Musculoskeletal Eval	o					o			o	o	o	o	o		
Pneumorax									•						
Psychosocial															
Development	s/o	s/o	s/o	s/o	s/o	s/o	s/o	s/o	s/o	s/o	s/o	s/o	s/o		
School Performance										o	o	o	o		
Socialization						s	s		s	s	s	s	s		

*Assure compliance with the AAP Recommendations for Preventive Pediatric Health Care

†Or at time of diagnosis

‡Discuss referral to specialist

§As needed

**Referral

|| Per state law

|| Once in this age group

#Every 2 years

††If hypercalciuria found, 2 repeat urine calcium (am and pm) should be sent. If still positive, repeat serum calcium, renal ultrasound for nephrocalcinosis and initiate dietary counseling

• = To be performed

S = Subjective (by history)

O = Objective (by a standard testing method)

Fig 2. Health supervision for children with Williams syndrome*.

should be confirmed by FISH analysis. The evaluations include the following:

- Complete physical and neurologic examination
- Growth parameters plotted on WS growth charts (Fig 3A–F)
- Cardiology evaluation
 - Full clinical evaluation by a cardiologist with expertise and experience in pediatric patients that includes 4-limb blood pressure measurements and echocardiography
- Genitourinary system evaluation
 - Ultrasonography of bladder and kidneys
 - Renal function studies (serum urea nitrogen and creatinine levels)
 - Urinalysis
- Calcium determinations (serum calcium, spot urine calcium, and creatinine levels) (Table 2)
- Thyroid function tests
- Ophthalmologic evaluation
- Multidisciplinary developmental evaluation (older than 2 years)
- FISH to determine ELN deletion

Referral to a clinical geneticist should be considered for individualized assessment and recommendations; a more extensive discussion of the clinical manifestations, natural history, recurrence risks, and future reproductive options; and evaluation of genetic risks for other family members.

SPECIAL CONSIDERATIONS FOR THE CHILD DIAGNOSED WITH WS

1. Do *not* give multivitamin preparations to children with WS because of the potential deleterious effects of vitamin D. Recommend diligent use of sunscreen to minimize autologous production of vitamin D.
2. Perform periodic cardiovascular evaluations, even after a baseline examination with normal findings.
3. Baseline cardiology evaluation should be performed by a cardiologist with pediatric expertise and experience.
4. Screen for the development of hypertension periodically according to guidelines of the American Academy of Pediatrics.
5. Establish a medical home with clear emphasis on continuity of care and the role of the family members as partners in the ongoing management and care of the child.

HEALTH SUPERVISION FROM BIRTH TO 1 YEAR (INFANCY)

Examination

1. Review and note clinical features and confirm diagnosis with FISH analysis
2. Routine health maintenance examinations and baseline evaluation
3. Growth and developmental evaluations using WS growth charts (Fig 3A–F)
4. Baseline cardiology evaluation by a cardiologist with pediatric expertise and experience
5. Review feeding issues (reflux, refusal, disordered suck or swallow, vomiting or symptoms of colic).

6. Consider pediatric ophthalmologic evaluation for strabismus, amblyopia, and refractive errors
7. Check for inguinal hernia
8. Objective hearing assessment at 6 to 12 months (recurrent otitis media is common)
9. Blood pressure measurement (both arms) annually and careful evaluation of femoral pulses
10. Early recognition and management of constipation
11. Pediatric anesthesia consultation for any child requiring surgery (several reports of unexpected deaths have been associated with the administration of anesthesia)²²

Laboratory

1. Williams Syndrome Chromosomal Region FISH to confirm clinical diagnosis
2. Serum creatinine level
3. Urinalysis
4. Calcium levels
 - a. Serum*
 - b. Spot urine test to determine calcium-creatinine ratio†
5. Thyroid screen for newborns (according to state mandate)
6. Baseline ultrasonographic examination of the bladder and kidneys

Anticipatory Guidance

1. Individual support for the family (by family, friends, clergy), support groups, or both (see list)
2. Review increased risk for otitis media
3. Feeding (difficulty in transition to textured foods)
4. Do *not* prescribe multivitamin preparations containing vitamin D
5. Refer to early childhood intervention program

HEALTH SUPERVISION FROM 1 TO 5 YEARS (EARLY CHILDHOOD)

Examination

1. Annual health maintenance examinations and baseline evaluation (including careful auscultation of chest and abdomen for murmurs or bruits)
2. Developmental evaluation and growth evaluation using WS growth charts (Fig 3A–F)
3. Annual cardiology evaluation from 1 to 5 years
4. Feeding issues: watch for rectal prolapse and avoid constipation with stool softeners if necessary
5. Annual hearing and vision screening; objective audiologic evaluation and an ophthalmologic evaluation before age 3 years
6. Orthopedic issues: musculoskeletal and neurologic assessments to evaluate joints, muscle tone, spasticity, and hyperactive reflexes¹⁷

*If hypercalcemia is found, dietary calcium restriction should be implemented and diet should be monitored in conjunction with a pediatric dietician/nutritionist. Referral to a pediatric renal specialist should be considered.

†If hypercalciuria is found, 2 repeated urine studies of the calcium-creatinine ratio (morning and afternoon) should be performed. If the level is still elevated, repeat measurement of the serum calcium level and perform renal ultrasonography for nephrocalcinosis. Assess dietary calcium intake.²¹




Fig 3A. Williams syndrome—stature, females.

7. Pediatric anesthesia consultation for any child requiring surgery (several reports of unexpected deaths have been associated with the administration of anesthesia)²²
8. Annual blood pressure measurement (both arms) and careful examination of femoral pulses
9. Multidisciplinary developmental assessment and treatment in early intervention programs (0–3 years) or school based programs (3 years and older)^{1,5,19}

10. Dental referral

Laboratory

1. Yearly urinalysis
2. Annual total calcium measurement if the level was elevated at baseline or as needed if the child becomes symptomatic; if level was normal, measure every 2 to 3 years
3. Urinary calcium-creatinine ratio every 2 years
4. Thyroid function test every 4 years

Height for males with Williams syndrome. Normal curves—dashed lines; affected patients—solid lines. Reprinted with permission from Saul RA, Geer JS, Seaver LH, Phelan MC, Sweet KM, Mills MS. *Growth References: Third Trimester to Adulthood*. Greenwood, SC: Greenwood Genetic Center; 1998.





Fig 3B. Williams syndrome—stature, males.

5. Serum creatinine level every 4 years

Anticipatory Guidance

1. Individual support for the family (by family, friends, clergy), support groups, or both
2. Review increased risk for otitis media
3. Ongoing feeding and dietary assessments

4. Therapy as needed (physical, speech and language, and occupational, including sensory integration)
5. Review constipation as a possible problem
6. Children with unexplained fever should be evaluated for urinary tract infection
7. Discuss developmental status, early intervention programs, and preschool programs



Weight for females with Williams syndrome. Normal curves—dashed lines; affected patients—solid lines. Reprinted with permission from Saul RA, Geer JS, Seaver LH, Phelan MC, Sweet KM, Mills MS. *Growth References: Third Trimester to Adulthood*. Greenwood, SC: Greenwood Genetic Center; 1998.

Fig 3C. Williams syndrome—weight, females.

HEALTH SUPERVISION FROM 5 YEARS TO 12 YEARS (LATE CHILDHOOD)

Examination

1. Annual health maintenance examinations and baseline evaluation
2. Developmental evaluation and growth evaluation using WS growth charts (Fig 3A–F)
3. Annual blood pressure measurements (both arms) and careful evaluation of femoral pulses
4. Cardiology evaluation as indicated by previous clinical findings. If results of previous evaluations are negative, repeated cardiology evaluation (for arterial stenoses, hypertension) should be performed at puberty
5. Ophthalmologic evaluation for strabismus and hyperopia
6. Orthopedic problems (eg, joint limitation, kyphosis, lordosis, scoliosis, and spasticity)

Weight for males with Williams syndrome. Normal curves—dashed lines; affected patients—solid lines. Reprinted with permission from Saul RA, Geer JS, Seaver LH, Phelan MC, Sweet KM, Mills MS. *Growth References: Third Trimester to Adulthood*. Greenwood, SC: Greenwood Genetic Center; 1998.




Fig 3D. Williams syndrome—weight, males.

7. Hearing and vision screening annually
8. Pediatric anesthesia consultation for any child requiring surgery (several reports of unexpected deaths have been associated with the administration of anesthesia²²)
9. School readiness and placement and Individual Educational Plan at 5 years
10. Developmental and psychoeducational assessment; formal evaluation for attention-deficit hyperactivity disorder, anxiety, or both and discussion of treatment options²³

Laboratory

1. Yearly urinalysis
2. Thyroid function tests every 4 years
3. Annual total calcium level if baseline result was elevated or child becomes symptomatic; otherwise measure level every 4 years
4. Urinary calcium-creatinine ratio every 2 years
5. Serum creatinine level every 2 to 4 years

Anticipatory Guidance

1. School readiness and placement

Head circumference for females with Williams syndrome. Normal curves—dashed lines; affected patients—solid lines. Reprinted with permission from Saul RA, Geer JS, Seaver LH, Phelan MC, Sweet KM, Mills MS. *Growth References: Third Trimester to Adulthood*. Greenwood, SC: Greenwood Genetic Center; 1998.




Fig 3E. Williams syndrome—head circumference, females.

2. Therapy as needed (physical, speech and language, and occupational, including sensory integration)
3. Long-term vocational planning
4. Discuss sexuality and adolescence; puberty is often early in WS, but true precocious puberty is rare
5. Discuss diet and exercise as obesity may become apparent in late childhood

6. Discuss treatment options for anxiety (counseling, relaxation techniques, and medications)
7. Estate planning for parents of a child with special needs

HEALTH SUPERVISION FROM 13 YEARS TO 18 YEARS (ADOLESCENCE)

Progressive medical problems including hypertension, progressive joint limitations, recurrent urinary




Fig 3F. Williams syndrome—head circumference, males.

tract infections, and gastrointestinal problems are common beginning in this age group and continuing throughout adult life.

Examination

1. Annual health maintenance examinations and baseline evaluation; blood pressure measurement (both arms)
2. Developmental evaluation and growth evaluation using WS growth charts (Fig 3A–F)
3. Cardiology evaluation if indicated by previous clinical findings
4. Pediatric anesthesia consultation for any child requiring surgery (several reports of unexpected deaths have been associated with the administration of anesthesia²²)
5. Consider ophthalmologic evaluation for hyperopia
6. Orthopedic problems (eg, joint limitation, kyphosis, lordosis, scoliosis, and spasticity)

TABLE 2. Normal Values for Random Urinary Calcium-Creatinine Ratios²¹

Age	Calcium-Creatinine Ratio (mg/mg ratio) (95th Percentile for Age)
<7 mo	0.86
7–18 mo	0.6
19 mo–6 y	0.42
Adults	0.22

7. Hearing and vision screening annually
8. Developmental and psychoeducational assessment; school placement and resource enhancement; vocational training; social skills training for peer interaction^{10,11}
9. Gastrointestinal issues: consider diverticulitis and diverticulosis, cholelithiasis, and chronic constipation in adolescents with abdominal pain
10. Screen for generalized anxiety disorder¹⁹

Laboratory

1. Yearly urinalysis
2. Thyroid function test every 4 years
3. Total calcium level only if adolescent becomes symptomatic, otherwise, every 4 years
4. Urinary calcium-creatinine ratio every 2 years
5. Bladder and renal ultrasonography at puberty and every 5 years thereafter
6. Serum creatinine level every 2 to 4 years

Anticipatory Guidance

1. School placement
2. Therapy as needed (physical, occupational, speech, and language)
3. Discuss diagnosis with the adolescent; support groups for the adolescent (see American Academy of Pediatrics statement on “Transition of Care Provided for Adolescents With Special Needs”)²⁴
4. Discuss sexuality and reproductive issues
5. Encourage career counseling
6. Foster independence
7. Assist in transition to adult care (especially for cardiology care). Many pediatricians feel comfortable continuing to provide primary care well into young adulthood
8. Encourage daily exercise to include range of motion
9. Encourage prompt medical attention for urinary tract or gastrointestinal symptoms
10. Mental health issues

COMMITTEE ON GENETICS, 2000–2001
 Christopher Cuniff, MD, Chairperson
 Jaime L. Frias, MD
 Celia I. Kaye, MD, PhD
 John Moeschler, MD
 Susan R. Panny, MD
 Tracy L. Trotter, MD

LIAISONS

Felix de la Cruz, MD, MPH
 National Institute of Child Health and Human Development

John Williams III, MD
 American College of Obstetricians and Gynecologists
 James W. Hanson, MD
 American College of Medical Genetics
 Cynthia A. Moore, MD, PhD
 Centers for Disease Control and Prevention
 Michele Lloyd-Puryear, MD, PhD
 Health Resources and Services Administration

SECTION LIAISON

H. Eugene Hoyme, MD
 Section on Genetics

CONSULTANTS

Paige Kaplan, MD
 Ron Lacro, MD
 Karen Levine, PhD
 Martin Levinson, MD
 Carolyn Mervis, PhD
 Colleen A. Morris, MD
 Beth A. Pletcher, MD
 Barbara Pober, MD
 Laurie Sadler, MD
 Paul Wang, MD

STAFF

Lauri A. Hall

REFERENCES

1. Williams JC, Barratt-Boyes BG, Lowe JB. Supravalvular aortic stenosis. *Circulation*. 1961;24:1311–1318
2. Beuren AJ. Supravalvular aortic stenosis: a complex syndrome with and without mental retardation. *Natl Found March Dimes Birth Defects Orig Art Ser*. 1972;8:45–56
3. Burn J. Williams syndrome. *J Med Genet*. 1986;23:389–395
4. Morris CA, Demsey SA, Leonard CO, Dilts C, Blackburn BL. Natural history of Williams syndrome: physical characteristics. *J Pediatr*. 1988;113:318–326
5. Martin ND, Yule W. A cognitive and behavioural phenotype in Williams syndrome. *J Clin Exp Neuropsychol*. 1991;13:232–244
6. Ewart AK, Morris CA, Atkinson D, et al. Hemizygosity at the elastin locus in a developmental disorder, Williams syndrome. *Nat Genet*. 1993;5:11–16
7. Lowery MC, Morris CA, Ewart A, et al. Strong correlation of elastin deletions, detected by FISH, with Williams syndrome: evaluation of 235 patients. *Am J Hum Genet*. 1995;57:49–53
8. Wu Y-Q, Sutton VR, Nickerson E, et al. Delineation of the common critical region in Williams syndrome and clinical correlation of growth, heart defects, ethnicity, and parental origin. *Am J Med Genet*. 1998;78:82–89
9. Martin ND, Snodgrass GJ, Cohen RD. Idiopathic infantile hypercalcemia: a continuing enigma. *Arch Dis Child*. 1984;59:605–613
10. Lopez-Rangel E, Maurice M, McGillivray B, Friedman JM. Williams syndrome in adults. *Am J Med Genet*. 1992;44:720–729
11. Morris CA, Leonard CO, Dilts C, Demsey SA. Adults with Williams syndrome. *Am J Med Gen Suppl*. 1990;6:102–107
12. Greenberg F, Lewis RA. The Williams syndrome: spectrum and significance of ocular features. *Ophthalmology*. 1988;95:1608–1612
13. Saul RA, Stevenson RE, Rogers RC, Skinner SA, Prouty LA, Flannery DB. Williams syndrome. In: *Proceedings of the Greenwood Genetic Center*. Greenwood, SC: Greenwood Genetic Center; 1988:204–209
14. Pankau R, Partsch C-J, Winter M, Gosch A, Wessel A. Incidence and spectrum of renal abnormalities in Williams-Beuren syndrome. *Am J Med Genet*. 1996;63:301–304
15. Pober BR, Lacro RV, Rice C, Mandell V, Teele RL. Renal findings in 40 individuals with Williams syndrome. *Am J Med Genet*. 1993;46:271–274
16. Morris CA, Mervis CB. Williams syndrome. In: Goldstein S, Reynolds CR, eds. *Handbook of Neurodevelopmental and Genetic Disorders in Children*. New York, NY: The Guilford Press; 1999:555–590
17. Kaplan P, Kirschner M, Watters G, Costa MT. Contractures in patients with Williams syndrome. *Pediatrics*. 1989;84:895–899

18. Wang PP, Hesselink JR, Jernigan TL, Doherty S, Bellugi U. Specific neurobehavioral profile of Williams' syndrome is associated with neocerebellar hemispheric preservation. *Neurology*. 1992;42:1999–2002
19. Chapman CA, du Plessis A, Pober BR. Neurologic findings in children and adults with Williams syndrome. *J Child Neurol*. 1996;11:63–65
20. Pober BR, Filiano JJ. Association of Chiari I malformations and Williams syndrome. *Pediatr Neurol*. 1995;12:84–88
21. Sargent JD, Stukel TA, Kresel J, Klein RZ. Normal values for random urinary calcium to creatinine ratios in infancy. *J Pediatr*. 1993;123:393–397
22. Bird LM, Billman GF, Lacro RV, et al. Sudden death in Williams syndrome: report of ten cases. *J Pediatr*. 1996;129:926–931
23. Power TJ, Blum NJ, Jones SM, Kaplan PE. Brief report: response to methylphenidate in two children with Williams syndrome. *J Autism Dev Dis*. 1997;27:79–87
24. American Academy of Pediatrics, Committee on Children With Disabilities. Transition of care provided for adolescents with special health care needs. *Pediatrics*. 1996;98:1203–1206

RESOURCES FOR PARENTS

March of Dimes, 1275 Mamaroneck Ave, White Plains, NY 10605; Telephone: 914/428–7100; <http://www.modimes.org>
The Williams Syndrome Association, PO Box 297, Clawson, MI 48017; Telephone: 248/541–3630; <http://www.williams-syndrome.org>
Williams Syndrome Foundation, University of California, Irvine, CA 92679; Telephone: 949/824–7259; <http://www.wsf.org>

Health Care Supervision for Children With Williams Syndrome

Committee on Genetics

Pediatrics 2001;107;1192-1204

Updated Information & Services	including high-resolution figures, can be found at: http://www.pediatrics.org/cgi/content/full/107/5/1192
References	This article cites 20 articles, 7 of which you can access for free at: http://www.pediatrics.org/cgi/content/full/107/5/1192#BIBL
Citations	This article has been cited by 4 HighWire-hosted articles: http://www.pediatrics.org/cgi/content/full/107/5/1192#otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Office Practice http://www.pediatrics.org/cgi/collection/office_practice
Errata	An erratum has been published regarding this article. Please see next page or: http://www.pediatrics.org/cgi/content/full/109/2/329
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.pediatrics.org/misc/Permissions.shtml
Reprints	Information about ordering reprints can be found online: http://www.pediatrics.org/misc/reprints.shtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



CONSULTANTS

Craig S. Derkay, MD
Seth M. Pransky, MD
Audie L. Whooley, MD

STAFF

Chelsea Kirk

BIBLIOGRAPHY

General Review

- Derkay CS, Caron JD, Wiatrak BJ, Choi SS, Jones JE. Postsurgical follow-up of children with tympanostomy tubes: results of the American Academy of Otolaryngology-Head and Neck Surgery Pediatric Otolaryngology Committee National Survey. *Arch Otolaryngol Head Neck Surg.* 2000;122:313-318
- Isaacson G, Rosenfeld RN. Care of the child with tympanostomy tubes: a visual guide for the pediatrician. *Pediatrics.* 1994;93:924-929

Audiologic Evaluation

- Emery M, Weber PC. Hearing loss due to myringotomy with tube placement and the role of preoperative audiograms. *Arch Otolaryngol Head Neck Surg.* 1998;124:421-424
- Manning SC, Brown OE, Roland PS, Phillips DZ. Incidence of sensorineural hearing loss in patients evaluated for tympanostomy tubes. *Arch Otolaryngol Head Neck Surg.* 1994;120:881-884

Otorrhea

- Chloe RA, Hubbell RN. Antimicrobial activity of silastic tympanostomy tubes impregnated with silver oxide. *Arch Otolaryngol Head Neck Surg.* 1995;121:562-565

- Force RW, Hart MC, Plummer SA, Powell DA, Nahata MC. Topical ciprofloxacin for otorrhea after tympanostomy tube placement. *Arch Otolaryngol Head Neck Surg.* 1995;121:880-884
- Mandel EM, Casselbrant ML, Kurs-lasky M. Acute otorrhea: bacteriology of a common complication of tympanostomy tubes. *Ann Otol Rhinol Laryngol.* 1994;103:713-718

Indications for Tympanostomy Tube Removal

- Cunningham MJ, Eavey RD, Krouse JH, Kiskaddon RM. Tympanostomy tubes: experience with removal. *Laryngoscope.* 1993;103:659-662

Sequelae of Tympanostomy Tube Extrusion/Removal

- Bulkley WJ, Bowes AK, Marlowe JF. Complications following ventilation of the middle ear using Goode T tubes. *Arch Otolaryngol Head Neck Surg.* 1991;117:895-898
- Matt BH, Miller RP, Meyers RM, Campbell JM, Cotton RT. Incidence of perforation with Goode T tubes. *Int J Pediatr Otorhinolaryngol.* 1991;21:1-6
- Maw AR, Bawden R. Tympanic membrane atrophy, scarring, atelectasis and attic retraction in persistent untreated OME and following ventilation tube insertion. *Int J Pediatr Otorhinolaryngol.* 1994;30:189-204
- Nichols PT, Ramadan HH, Wax MK, Santrock RD. Relationship between tympanic membrane perforation and retained ventilation tubes. *Arch Otolaryngol Head Neck Surg.* 1998;124:417-419
- Tos M, Stangerup S-E. Hearing loss in tympanosclerosis caused by grommets. *Arch Otolaryngol Head Neck Surg.* 1989;115:931-935
- Weigel MT, Parker MY, Goldsmith MM, Postma DS, Pillsbury HC. A prospective randomized study of four commonly used tympanostomy tubes. *Laryngoscope.* 1989;99:252-256

ERRATUM

Several errors occurred in a figure in the policy statement "Health Care Supervision for Children With Williams Syndrome," which appeared in the May 2001 issue of *Pediatrics* (2001;107:1192-1204). In Fig 2, first column, the 11th row heading under Medical Evaluation should read "Musculoskeletal Eval," and the 12th row heading should read "Pneumovax." In the footnotes, the explanation with the double dagger should read "If hypercalciuria is found, 2 repeated urine studies of the calcium-creatinine ratio (morning and afternoon) should be performed. If the level is still elevated, repeat measurement of the serum calcium level and perform renal ultrasonography for nephrocalcinosis. Assess dietary calcium intake.²¹" The explanation for the abbreviation O should read "Objective . . ."